



The Faber Polynomial Expansion Method for a Subclass of Analytic and Bi-Univalent Functions Associated with the Janowski Functions

Nazar Khan ^a, Qazi Zahoor Ahmad ^b, Shahid Khan^c, Bilal Khan ^d

^aDepartment of Mathematics, Abbottabad University of Science and Technology, Abbottabad 22010, Pakistan

^bDepartment of Mathematics, Abbottabad University of Science and Technology, Abbottabad 22010, Pakistan

^cDepartment of Mathematics Riphah International University Islamabad, Pakistan

^dSchool of Mathematical Sciences, East China Normal University, 500 Dongchuan Road, Shanghai 200241, People's Republic of China

Abstract

In this present investigation, we introduce a new subclass of analytic and bi-univalent functions associated with Janowski functions. Using the Faber polynomial expansions, we determine a general coefficients bounds $|a_n|$, $n \geq 3$ for this newly defined class. Relevant connections of the results presented in this paper with those in a number of other related works on this subject are also pointed out.

Keywords: Analytic functions, Univalent functions, Bi-univalent functions, Faber polynomial expansion

2010 MSC: 05A30, 30C45, 11B65, 47B38

1. Introduction and Definitions

Let $\mathcal{H}(\mathbb{U})$ denote the class of functions analytic in the open unit disc

$$\mathbb{U} = \{z : z \in \mathbb{C}, |z| < 1\}$$

and let \mathcal{A} be the subclass of functions f in $\mathcal{H}(\mathbb{U})$ satisfying the normalization condition

$$f(0) = f'(0) - 1 = 0.$$

Thus, the functions in \mathcal{A} are represented by the Taylor-Maclaurin series expansion

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n \quad (z \in \mathbb{U}). \quad (1.1)$$

Let \mathcal{S} be the class of univalent functions in \mathcal{A} . Furthermore, let \mathcal{P} denotes the class of analytic functions ϕ normalized by

$$\phi(z) = 1 + \sum_{n=1}^{\infty} p_n z^n \quad (z \in \mathbb{U}), \quad (1.2)$$

†Article ID: MTJPAM-D-20-00006

Email addresses: nazarmaths@gmail.com (Nazar Khan ) , zahoorqazi5@gmail.com (Qazi Zahoor Ahmad ) , shahidmath761@gmail.com (Shahid Khan), bilalmaths789@gmail.com (Bilal Khan )

Received:27 March 2020, Accepted:1 December 2020, Published:25 April 2021

*Corresponding Author: Nazar Khan



such that

$$\Re(\phi(z)) > 0.$$

Definition 1.1. A function $f \in \mathcal{A}$ is said to be subordinate to a function g written as $f < g$, if there exist a Schwarz function w with

$$w(0) = 0 \quad \text{and} \quad |w(z)| < 1 \quad \text{such that} \quad f = g(w).$$

In particular if g is univalent in \mathbb{U} , then

$$f(0) = g(0) \quad \text{and} \quad f(\mathbb{U}) \subset g(\mathbb{U}).$$

Definition 1.2. A function p is said to be in the class $\mathcal{P}[A, B]$ ($-1 \leq B < A \leq 1$) if it is analytic in \mathbb{U} with $p(0) = 1$ and

$$p(z) < \frac{1 + Az}{1 + Bz}, \quad (z \in \mathbb{U}).$$

Geometrically a function p belongs to $\mathcal{P}[A, B]$ maps the open unit disc \mathbb{U} onto the disc defined by the domain

$$\Omega[A, B] = \left\{ w : \left| w - \frac{1 - AB}{1 - B^2} \right| < \frac{A - B}{1 - B^2} \right\}.$$

The class $\mathcal{P}[A, B]$ connected with the class \mathcal{P} of functions with positive real part by the relation

$$p(z) \in \mathcal{P}, \quad \text{if and only if} \quad \frac{(A + 1)p(z) - (A - 1)}{(B + 1)p(z) - (B - 1)} \in \mathcal{P}[A, B].$$

The class $\mathcal{P}[A, B]$ was introduced by Janowski [10]. This functions class is further generalized by the many well-known authors (see, for example [11, 15, 23, 19, 24]).

The Koebe one-quarter theorem shows that the image of \mathbb{U} under every univalent function $f \in \mathcal{A}$ contains a disk $\{w : |w| < \frac{1}{4}\}$ of radius $\frac{1}{4}$. Every univalent function f has an inverse f^{-1} defined on some disk containing the disk $\{w : |w| < \frac{1}{4}\}$ and satisfying:

$$f^{-1}(f(z)) = z \quad (z \in \mathbb{U}),$$

and

$$f(f^{-1}(w)) = w \quad \left(|w| < r_0(f), \quad r_0(f) \geq \frac{1}{4} \right),$$

where

$$f^{-1}(w) = w - a_2w^2 + (2a_2^2 - a_3)w^3 - (5a_2^3 - 5a_2a_3 + a_4)w^4 + \dots \tag{1.3}$$

A function $f \in \mathcal{S}$ is said to be bi-univalent on \mathbb{U} if f and $g = f^{-1}$ are both univalent on \mathbb{U} . We denote the class of all such functions by Σ . In recent years, the pioneering work of Srivastava *et al.* [20] essentially revived the investigation of various subclasses of the analytic and bi-univalent functions class Σ . In fact, in a remarkably large number of sequels to the pioneering work of Srivastava *et al.* [20], several different subclasses of the analytic and bi-univalent functions class Σ were introduced and studied analogously by the many authors (see, for example [4, 5, 6, 17, 12, 27, 26, 28]). However, only non-sharp estimates on the initial coefficients $|a_2|$ and $|a_3|$ in the Taylor-Maclaurin expansion (1.1) were obtained in these recent papers.

The Faber polynomials introduced by Faber [8] play an important role in various areas of mathematical sciences, especially in Geometric Function Theory of Complex Analysis. Recently, several authors (see, for example [9, 21, 25, 16, 18]) investigated some interesting and useful properties for analytic functions by applying the Faber polynomial expansion method. Motivated by these and other recent work here we define new subclasses of analytic and bi-univalent functions in \mathbb{U} and (by means of the Faber polynomial expansion method) we determine estimates for the general coefficients $|a_n|$ ($n \geq 3$) in the Taylor-Maclaurin series expansion (1.1) of functions in each of these subclasses.

We now introduce the following subclass of the analytic and bi-univalent functions class Σ .

Definition 1.3. Let $b \in \mathbb{C} \setminus \{0\}$, $\beta \geq 0$, $\lambda \geq 0$ and $-1 \leq B < A \leq 1$. A function $f \in \mathcal{A}$ is said to belong to the class $\mathcal{R}[b, \beta, \lambda, A, B]$, if and only if

$$1 + \frac{1}{b} \{\mathcal{L}(f, \beta, \lambda; z) - 1\} < \frac{1 + Az}{1 + Bz}, \quad z \in \mathbb{U}, \tag{1.4}$$

and

$$1 + \frac{1}{b} \{\mathcal{M}(g, \beta, \lambda; w) - 1\} < \frac{1 + Aw}{1 + Bw}, \quad w \in \mathbb{U}. \tag{1.5}$$

where

$$\begin{aligned} \mathcal{L}(f, \beta, \lambda; z) &= \frac{(1 - \beta) f(z) + \beta (zf'(z)) + \lambda z^2 f''(z)}{z}, \\ \mathcal{M}(g, \beta, \lambda; w) &= \frac{(1 - \beta) g(w) + \beta (wg'(w) + \lambda w^2 g''(w))}{w} \end{aligned}$$

and $g(w) = f^{-1}(w)$ defined in (1.3).

2. The Faber Polynomial Expansion Method and Its Applications

In this section, by using the Faber polynomial expansion of a function f of the form (1.1), we observe that the coefficients of its inverse map $g = f^{-1}$ may be expressed as follows;

$$g(w) = f^{-1}(w) = w + \sum_{n=2}^{\infty} \frac{1}{n} K_{n-1}^{-n}(a_2, a_3, \dots) w^n,$$

where

$$\begin{aligned} K_{n-1}^{-n} &= \frac{(-n)!}{(-2n+1)!(n-5)!} a_2^{n-1} + \frac{(-n)!}{[2(-n+1)]!(n-3)!} a_2^{n-3} a_3 + \frac{(-n)!}{(-2n+3)!(n-4)!} a_2^{n-4} a_4 \\ &+ \frac{(-n)!}{[2(-n+2)]!(n-5)!} a_2^{n-5} [a_5 + (-n+2)a_3^2] + \frac{(-n)!}{(-2n+5)!(n-6)!} a_2^{n-6} [a_6 + (-2n+5)a_3 a_4] + \sum_{j \geq 7} a_2^{n-j} V_j, \end{aligned}$$

Here, and in what follows, such expressions as (for example) $(-n)!$ are to be interpreted symbolically by

$$(-n)! = \Gamma(1-n) = (-n)(-n-1)(-n-2) \dots$$

and V_j ($7 \leq j \leq n$) is a homogeneous polynomial in the variables a_2, a_3, \dots, a_n [3]. In particular, the first three terms of K_{n-1}^{-n} are given below

$$\frac{1}{2} K_1^{-2} = -a_2, \tag{2.1}$$

$$\frac{1}{3} K_2^{-3} = 2a_2^2 - a_3 \tag{2.2}$$

and

$$\frac{1}{4} K_3^{-4} = -(5a_2^3 - 5a_2 a_3 + a_4). \tag{2.3}$$

In general, an expansion of K_n^p is given by (see, for details, [2]),

$$K_n^p = p a_n + \frac{p(p-1)}{2} E_{n-1}^2 + \frac{p!}{(p-3)!3!} E_{n-1}^3 + \dots + \frac{p!}{(p-n+1)!(n-1)!} E_{n-1}^{n-1} \quad p \in \mathbb{Z}, \tag{2.4}$$

where $\mathbb{Z} = \{0, \pm 1, \pm 2, \dots\}$ (see [13]).

$$E_n^p = E_n^p(a_2, a_3, \dots).$$

It is clearly seen that

$$E_{n-1}^m(a_2, \dots, a_n) = \sum_{n=2}^{\infty} \frac{m!(a_2)^{\mu_1} \dots (a_n)^{\mu_{n-1}}}{\mu_1! \dots \mu_{n-1}!} \quad (m \leq n).$$

While $a_1 = 1$, and the sum is taken over all nonnegative integer μ_1, \dots, μ_n satisfying:

$$\mu_1 + \mu_2 + \dots + \mu_n = m,$$

and

$$\mu_1 + 2\mu_2 + \dots + (n-1)\mu_{n-1} = n-1.$$

Evidently, (see [1])

$$E_{n-1}^{n-1}(a_2, \dots, a_n) = a_2^{n-1},$$

or equivalently,

$$E_n^m(a_1, a_2, \dots, a_n) = \sum_{n=1}^{\infty} \frac{m!(a_1)^{\mu_1} \dots (a_n)^{\mu_n}}{\mu_1! \dots \mu_n!}, \quad m \leq n,$$

again $a_1 = 1$, and taking the sum over all nonnegative integer μ_1, \dots, μ_n satisfying:

$$\mu_1 + \mu_2 + \dots + \mu_n = m,$$

and

$$\mu_1 + 2\mu_2 + \dots + (n)\mu_n = n.$$

It is clear that

$$E_n^n(a_1, \dots, a_n) = E_1^n$$

the first and last polynomials are

$$E_n^n = a_1^n \quad \text{and} \quad E_n^1 = a_n.$$

Now before to state and prove our main results we need the following lemma.

Lemma 2.1. [9] Let the function $w \in \mathcal{A}$ given by

$$w(z) = \sum_{n=1}^{\infty} w_n z^n$$

be a Schwarz function so that

$$|w(z)| < 1 \quad \text{for} \quad |z| < 1.$$

If $\gamma \geq 0$ then

$$|w_2 + \gamma w_1^2| \leq 1 + (\gamma - 1)|w_1|^2.$$

Theorem 2.2. For $b \in \mathbb{C} \setminus \{0\}$, $\beta \geq 0$, $\lambda \geq 0$, and $-1 \leq B < A \leq 1$. Let $f \in \mathcal{R}[b, \beta, \lambda, A, B]$, if $a_m = 0$, $2 \leq m \leq n-1$, then

$$|a_n| \leq \frac{|b|(A-B)}{1+(n-1)(\beta+n\lambda)} \quad (n \geq 3). \tag{2.5}$$

Proof. For the function $f \in \mathcal{R}[b, \beta, \lambda, A, B]$ of the form (1.1), we have

$$1 + \frac{1}{b} \{ \mathcal{L}(f, \beta, \lambda; z) - 1 \} = 1 + \sum_{n=2}^{\infty} \frac{1+(n-1)(\beta+n\lambda)}{b} a_n z^{n-1}, \tag{2.6}$$

and for its inverse map $g = f^{-1}$, we have

$$1 + \frac{1}{b} \{ \mathcal{M}(g, \beta, \lambda; w) - 1 \} = 1 + \sum_{n=2}^{\infty} \frac{1 + (n-1)(\beta + n\lambda)}{b} A_n w^{n-1}, \tag{2.7}$$

where,

$$A_n = \frac{1}{n} K_{n-1}^{-n}(a_2, a_3, \dots, a_n).$$

Since, both functions f and its inverse $g = f^{-1}$ are in $\mathcal{R}[b, \beta, \lambda, A, B]$, by the definition of subordination, there exist two Schwarz functions

$$u(z) = \sum_{n=1}^{\infty} c_n z^n \quad (z \in \mathbb{U})$$

and

$$v(w) = \sum_{n=1}^{\infty} d_n w^n \quad (w \in \mathbb{U}).$$

So that we have

$$\begin{aligned} 1 + \frac{1}{b} \{ \mathcal{L}(f, \beta, \lambda; z) - 1 \} &= \frac{1 + A(u(z))}{1 + B(u(z))} \\ &= 1 - \sum_{n=1}^{\infty} (A - B) K_n^{-1}(c_1, c_2, \dots, c_n, B) z^n \end{aligned} \tag{2.8}$$

and

$$\begin{aligned} 1 + \frac{1}{b} \{ \mathcal{M}(g, \beta, \lambda; w) - 1 \} &= \frac{1 + A(v(w))}{1 + B(v(w))} \\ &= 1 - \sum_{n=1}^{\infty} (A - B) K_n^{-1}(d_1, d_2, \dots, d_n, B) w^n. \end{aligned} \tag{2.9}$$

In general (see, for example [2, 3]) for any $p \in \mathbb{N}$ and $n \geq 2$, an expansion of $K_n^p(k_1, k_2, \dots, k_n, B)$,

$$\begin{aligned} K_n^p(k_1, k_2, \dots, k_n, B) &= \frac{p!}{(p-n)!n!} k_1^n B^{n-1} + \frac{p!}{(p-n+1)!(n-2)!} k_1^{n-2} k_2 B^{n-2} \\ &+ \frac{p!}{(p-n+2)!(n-3)!} \times k_1^{n-3} k_3 B^{n-3} \\ &+ \frac{p!}{(p-n+3)!(n-4)!} k_1^{n-4} \left[k_4 B^{n-4} + \frac{p-n+3}{2} k_3^2 B \right] \\ &+ \frac{p!}{(p-n+4)!(n-5)!} k_1^{n-5} \left[k_5 B^{n-5} + (p-n+4) k_3 k_4 B \right] + \sum_{j \geq 6} k_1^{n-1} X_j, \end{aligned}$$

where X_j is a homogeneous polynomial of degree j in the variables k_1, k_2, \dots, k_n .

For the coefficients of the Schwarz functions $u(z)$ and $v(w)$ we have (see, for example [7])

$$|c_n| \leq 1 \quad \text{and} \quad |d_n| \leq 1.$$

Comparing the corresponding coefficients of (2.6) and (2.8), we have

$$\frac{1 + (n-1)(\beta + n\lambda)}{b} a_n = -(A - B) K_{n-1}^{-1}(c_1, c_2, \dots, c_{n-1}, B), \tag{2.10}$$

Similarly corresponding coefficients of (2.7) and (2.9), we have

$$\frac{1 + (n-1)(\beta + n\lambda)}{b} A_n = -(A - B) K_n^{-1}(d_1, d_2, \dots, d_n, B) \tag{2.11}$$

Note that for $a_m = 0; 2 \leq m \leq n - 1$, we have $A_n = -a_n$

$$\frac{1 + (n - 1)(\beta + n\lambda)}{b} a_n = -(A - B)c_{n-1}, \tag{2.12}$$

$$\frac{1 + (n - 1)(\beta + n\lambda)}{b} a_n = (A - B)d_{n-1}. \tag{2.13}$$

Taking the absolute values of (2.12) and (2.13), we have

$$|a_n| \leq \frac{b(A - B)}{1 + (n - 1)(\beta + n\lambda)} |c_{n-1}| = \frac{b(A - B)}{1 + (n - 1)(\beta + n\lambda)} |d_{n-1}|,$$

$$|a_n| \leq \frac{b(A - B)}{1 + (n - 1)(\beta + n\lambda)}, \quad n \geq 3.$$

which completes the proof of Theorem 2.2. □

If we set

$$\beta = b = 1 = -B = A + 2\alpha \quad (0 \leq \alpha < 1)$$

in Theorem 2.2, we obtain the following known result.

Corollary 2.3. [22] Let $f \in \mathcal{R}[1, 1, \lambda, 1 - 2\alpha, -1]$, if $a_m = 0, 2 \leq m \leq n - 1$, then

$$|a_n| \leq \frac{2(1 - \alpha)}{n[1 + \lambda(n - 1)]}, \quad n \geq 3.$$

Theorem 2.4. For $b \in \mathbb{C} \setminus \{0\}, \beta \geq 0, \lambda \geq 0$ and $-1 \leq B < A \leq 1$. Let $f \in \mathcal{R}[b, \beta, \lambda, A, B]$ then

$$|a_2| \leq \begin{cases} \frac{b(A-B)}{\sqrt{b(A-B)\{1+2(\beta+3\lambda)\}+(1+B)\{1+(\beta+2\lambda)\}^2}}, & \text{if } 0 \leq B < A, \\ \frac{b(A-B)}{1+(\beta+2\lambda)}, & \text{otherwise.} \end{cases}$$

and

$$|a_3 - 2a_2^2| \leq \frac{b^2(A-B)^2 - (B+1)\{1+\beta+2\lambda\}^2}{b(A-B)[1+2(\beta+3\lambda)]} |a_2|^2, \quad \text{if } A \leq 0,$$

$$|a_3 - a_2^2| \leq \frac{b(A-B)}{(1+2(\beta+3\lambda))}, \quad \text{if } A > 0.$$

Proof. Replacing n by 2 and 3 in (2.10) and (2.11), respectively, we have

$$\frac{1 + (\beta + 2\lambda)}{b} a_2 = (A - B)c_1, \tag{2.14}$$

$$\frac{1 + 2(\beta + 3\lambda)}{b} a_3 = (A - B)(Bc_1^2 - c_2), \tag{2.15}$$

$$- \frac{1 + (\beta + 2\lambda)}{b} a_2 = (A - B)d_1, \tag{2.16}$$

$$\frac{1 + 2(\beta + 3\lambda)}{b} \{2a_2^2 - a_3\} = (A - B)(Bd_1^2 - d_2). \tag{2.17}$$

From (2.14) and (2.16) we have

$$|a_2| \leq \frac{b(A - B)}{1 + (\beta + 2\lambda)} |c_1| = \frac{b(A - B)}{1 + (\beta + 2\lambda)} |d_1|,$$

$$\leq \frac{b(A - B)}{1 + (\beta + 2\lambda)}.$$
(2.18)

Adding (2.15) and (2.17) we have

$$\frac{2(1+2(\beta+3\lambda))}{b} a_2^2 = (A-B) \{ (Bc_1^2 - c_2) + (Bd_1^2 - d_2) \}. \tag{2.19}$$

Taking absolute values of (2.19), we obtain

$$\frac{2(1+2(\beta+3\lambda))}{b} |a_2|^2 = (A-B) \{ |c_2 + (-B)c_1^2| + |d_2 + (-B)d_1^2| \}$$

If $B \leq 0$, then by Lemma 2.1, we have

$$\frac{2(1+2(\beta+3\lambda))}{b} |a_2|^2 \leq (A-B) \{ 1 + (-B-1)|c_1|^2 + 1 + (-B-1)|d_1|^2 \}. \tag{2.20}$$

By using

$$\left(\frac{1+(\beta+2\lambda)}{b(A-B)} \right)^2 |a_2|^2 = |c_1|^2 = |d_1|^2,$$

we can get that

$$|a_2| \leq \frac{b(A-B)}{\sqrt{b(A-B)\{1+2(\beta+3\lambda)\} + (1+B)\{1+(\beta+2\lambda)\}^2}}.$$

Obviously, for $A > 0$ we have

$$\frac{b(A-B)}{\sqrt{b(A-B)\{1+2(\beta+3\lambda)\} + (1+B)\{1+(\beta+2\lambda)\}^2}} < \frac{b(A-B)}{1+(\beta+2\lambda)}.$$

Rewriting equation (2.17), we have

$$\frac{1+2(\beta+3\lambda)}{b} \{ a_3 - 2a_2^2 \} = (A-B) \{ d_2 - Bd_1^2 \}.$$

Taking the absolute values of the above equation, we have

$$\left| \frac{1+2(\beta+3\lambda)}{b} \{ a_3 - 2a_2^2 \} \right| = (A-B) \left| \{ d_2 - Bd_1^2 \} \right|.$$

we can get that If $A \leq 0$, then by Lemma 2.1 and

$$a_2^2 = \left(\frac{b(A-B)}{1+(\beta+2\lambda)} \right)^2 d_1^2,$$

we have

$$|a_3 - 2a_2^2| \leq \frac{b^2(A-B)^2 - (B+1)\{1+\beta+2\lambda\}^2}{b(A-B)[1+2(\beta+3\lambda)]} |a_2|^2.$$

By using

$$\left(\frac{1+(\beta+2\lambda)}{b(A-B)} \right)^2 |a_2|^2 = |d_1|^2,$$

for $A > 0$, subtracting (2.17) from (2.15), we have

$$\frac{2(1+2(\beta+3\lambda))}{b} (a_3 - a_2^2) = (A-B) [B(d_1^2 - c_1^2) + (c_2 - d_2)].$$

Using the fact that $c_1^2 = d_1^2$ and taking the absolute values of both sides of the above equation, we obtain the desired inequality

$$|a_3 - a_2^2| \leq \frac{b(A-B)}{(1+2(\beta+3\lambda))}.$$

This completes the proof. □

Acknowledgments

This paper is dedicated to Professor Hari Mohan Srivastava on the occasion of his 80th Birthday.

References

- [1] H. Airault, *Symmetric sums associated to the factorizations of Grunsky coefficients*, in Conference, Groups and Symmetries. Montreal, Canada, 2007.
- [2] H. Airault, *Remarks on Faber polynomials*, Int. Math. Forum. **3** (9), 449-456, 2008.
- [3] H. Airault and H. Bouali, *Differential calculus on the Faber polynomials*, Bull. Sci. Math. **130** (3), 179-222, 2006.
- [4] Ş. Altınkaya and S. Yalçın, *Faber polynomial coefficient bounds for a subclass of bi-univalent functions*, C. R. Acad. Sci. Paris, Ser. I **353**, 1075-1080, 2015.
- [5] D. A. Brannan and T. S. Taha, *On some classes of bi-univalent functions*, in: S.M. Mazhar, A. Hamoui, N.S. Faour (Eds.), Mathematical Analysis and its Applications, Kuwait, 1985, in: KFAS Proceedings Series, vol. 3, Pergamon Press, Elsevier Science Limited, Oxford, 1988, pp. 53-60; see also Studia Univ. Babeş-Bolyai Math. **31** (2), 70-77, 1986.
- [6] S. Bulut, *Faber polynomial coefficient estimates for a comprehensive subclass of analytic bi-univalent functions*, C. R. Acad. Sci. Paris, Ser. I **352**, 479-484, 2014.
- [7] P. L. Duren, *Univalent Functions*, Grundlehren der Math. Wiss., Vol. 259, Springer-Verlag, New York, 1983.
- [8] G. Faber, *Über polynomische Entwicklungen*, Math. Ann. **57** (3), 1569-1573, 1903.
- [9] S. G. Hamidi and J. M. Jahangiri, *Faber polynomial coefficients of bi-subordinate functions*, Comptes Rendus Mathematique. **354**, 365-370, 2016.
- [10] W. Janowski, *Some extremal problems for certain families of analytic functions*, Ann. Polon. Math. **28**, 297-326, 1973.
- [11] S. Mahmood, Q. Z. Ahmad, H. M. Srivastava, N. Khan, B. Khan and M. Tahir, *A certain subclass of meromorphically q -starlike functions associated with the Janowski functions*, J. Inequal. Appl. **2019** (88), 1-11, 2019.
- [12] M. Sabil, Q. Z. Ahmad, B. Khan M. Tahir and N. Khan, *Generalisation of certain subclasses of analytic and univalent functions*, Maejo Internat. J. Sci. Technol. **13** (01), 1-9, 2019.
- [13] A. C. Schiffer and D. C. Spencer, *The coefficient of Schlicht functions*, Duke Math. J. **10**, 611-635, 1943.
- [14] M. Schiffer, *A method of variation within the family of simple functions*, Proc. Lond. Math. Soc. **44** (2), 432-449, 1938.
- [15] H. M. Srivastava, Q. Z. Ahmad, N. Khan, N. Khan and B. Khan, *Hankel and Toeplitz determinants for a subclass of q -starlike functions associated with a general conic domain*, Mathematics **7** (2), 181, 1-15, 2019.
- [16] H. M. Srivastava, Ş. Altınkaya and S. Yalçın, *Certain subclasses of bi-univalent functions associated with the Horadam polynomials*, Iran. J. Sci. Technol. Trans. A: Sci. **43**, 1873-1879, 2019.
- [17] H. M. Srivastava, S. Bulut, M. Caglar and N. Yagmur, *Coefficient estimates for a general subclass of analytic and bi-univalent functions*, Filomat **27**, 831-842, 2013.
- [18] H. M. Srivastava, S. Khan, Q. Z. Ahmad, N. Khan and S. Hussain, *The Faber polynomial expansion method and its application to the general coefficient problem for some subclasses of bi-univalent functions associated with a certain q -integral operator*, Stud. Univ. Babeş-Bolyai Math. **63**, 419-436, 2018.
- [19] H. M. Srivastava, B. Khan, N. Khan and Q. Z. Ahmad, *Coefficient inequalities for q -starlike functions associated with the Janowski functions*, Hokkaido Math. J. **48**, 407-425, 2019.
- [20] H. M. Srivastava, A. K. Mishra and P. Gochhayat, *Certain subclasses of analytic and bi-univalent functions*, Appl. Math. Lett. **23**, 1188-1192, 2010.
- [21] H. M. Srivastava, A. Motamednezhad and E. A. Adegan, *Faber polynomial coefficient estimates for bi-univalent functions defined by using differential subordination and a certain fractional derivative operator*, Mathematics **8** (2), 172, 1-12, 2020.
- [22] H. M. Srivastava, S. Sümer Eker and R. M. Ali, *Coefficient bounds for a certain class of analytic and bi-univalent functions*, Filomat **29** (8), 1839-1845, 2015.
- [23] H. M. Srivastava, M. Tahir, B. Khan, Q. Z. Ahmad and N. Khan, *Some general classes of q -starlike functions associated with the Janowski functions*, Symmetry **11** (2), 292, 1-14, 2019.
- [24] H. M. Srivastava, M. Tahir, B. Khan, Q. Z. Ahmad and N. Khan, *Some general families of q -starlike functions associated with the Janowski functions*, Filomat **33** (9), 2613-2626, 2019.
- [25] H. M. Srivastava and A. K. Wanas, *Initial Maclaurin coefficient bounds for new subclasses of analytic and m -fold symmetric bi-univalent functions defined by a linear combination*, Kyungpook Math. J. **59**, 493-503, 2019.
- [26] T. S. Taha, *Topics in Univalent Function Theory*, Ph.D. Thesis, University of London, 1981.
- [27] M. Tahir, N. Khan, Q. Z. Ahmad, B. Khan and G. Mehtab, *Coefficient estimates for some subclasses of analytic and bi-univalent functions associated with conic domain*, Sahand Communications in Math. Anal. (SCMA) **16** (1), 69-81, 2019.
- [28] Q.-H. Xu, H.-G. Xiao and H. M. Srivastava, *A certain general subclass of analytic and bi-univalent functions and associated coefficient estimate problems*, Appl. Math. Comput. **218**, 11461-11465, 2012.