

On generalizations of Tribonacci numbers

Ortaç Öneş ^a, Mustafa Alkan ^b

^aAkdeniz University, Department of Mathematics, Antalya 07058, Turkey

^bAkdeniz University, Department of Mathematics, Antalya 07058, Turkey

Abstract

In this article, some generalizations of the results in the literature are obtained by using the sequences over an integral domain with the help of matrix method. Then some generalizations for the d'Ocagne identity, the Honsberger's formula, the Cassini's identity, the Catalan's identity are given. Finally, Binet formulas of sequences in the literature are unified in a theorem.

Keywords: Integral domain, Tribonacci numbers, Tribonacci sequences

2010 MSC: 05A19, 11B39

1. Introduction

After being defined the Fibonacci numbers in 13th century, it has been in the center of interest of most of researchers. Then a lot of researchers have generalized the Fibonacci numbers to the Fibonacci polynomials, the Pell numbers and polynomials, the Lucas numbers and polynomials, the Jacobstal numbers and polynomials, the Pell-Lucas numbers and polynomials, the Jacobstal-Lucas numbers and polynomials, the Chebyshev numbers and polynomials, which are derived from the Fibonacci numbers ([2], [3], [5], [7], [16], [17], [19], [24], [25], [27]).

In the ring theory, it is well-known that the polynomial rings with finite indeterminates over integral domains are an integral domain. Thus in this article, we deal with the generalizations of the Tribonacci sequences over an integral domain and examine them with the help of the matrix method.

In 1965, Horadam defined the sequences of numbers, which are called as Horadam numbers, on rational numbers in [7] as follows:

$$h_n = ph_{n-1} - qh_{n-2} \quad (n \geq 3)$$



with the initial conditions $h_1 = a$ and $h_2 = bx$.

The Tribonacci numbers in [12] was defined as $T_n = T_{n-1} + T_{n-2} + T_{n-3}$ for $n \geq 4$ with the initial conditions $T_1 = T_2 = 1$ and $T_3 = 2$ and the Tribonacci numbers were generalized to the Tribonacci polynomials in [22].

Definition 1.1 (cf. [18]). Let p, q, r, a and b be elements of an integral domain R . Then, for a positive integer $n \geq 2$, the recurrence relations are defined as

$$w_{n+1} = pw_n + qw_{n-1} + rw_{n-2},$$

†Article ID: MTJPAM-D-21-00076

Email addresses: ortacs@akdeniz.edu.tr (Ortaç Öneş ) , alkan@akdeniz.edu.tr (Mustafa Alkan )

Received:17 December 2021, Accepted:7 January 2022

*Corresponding Author: Ortaç Öneş

where $w_0 = a, w_1 = b, w_2 = pw_1 + qw_0$ and it is said that

$$w = \{w_{n+1} = pw_n + qw_{n-1} + rw_{n-2} : n \in \mathbb{Z}^+, n \geq 2\}$$

is the sequence of p, q, r with a, b on R denoted by $w_n(a, b, p, q, r)$ or briefly $\{w_n\}$.

It is clear that Definition 1.1 generalizes not only the Horadam numbers and polynomials in [7], but also the Tribonacci numbers and polynomials, the Tricobstal numbers and polynomials. Moreover, when $p \neq 0, q \neq 0$ and $r = 0$, then Definition 1.1 is equivalent to the definition in [1].

Now, we fix the notations in Definition 1.1.

Remark 1.2. Let $R = \mathbb{Z}[x, y]$. If $p = x, q = y, r = 0, a = 0$ and $b = 1$, then $w_n(x, y)$ is the n -th Fibonacci polynomial with two variables for positive integer n .

Fibonacci numbers	Fibonacci polynomials with two variables
0	$w_0(x, y) = 0$
1	$w_1(x, y) = 1$
1	$w_2(x, y) = x$
2	$w_3(x, y) = x^2 + y$
3	$w_4(x, y) = x^3 + 2xy$
5	$w_5(x, y) = x^4 + 3x^2y + y^2$
8	$w_6(x, y) = x^5 + 4x^3y + 3xy^2$
13	$w_7(x, y) = x^6 + 5x^4y + 6x^2y^2 + y^3$

Example 1.3. Let $p = x^2, q = x$ and $r = 1$ be elements of an integral domain $R = \mathbb{Z}[x]$. Then we recall that $w_n(x)$ is called Tribonacci polynomials when $a = 0$ and $b = 1$.

Tribonacci numbers	Tribonacci polynomials
0	$w_0(x) = 0$
1	$w_1(x) = 1$
1	$w_2(x) = x^2$
2	$w_3(x) = x^4 + x$
4	$w_4(x) = x^6 + 2x^3 + 1$
7	$w_5(x) = x^8 + 3x^5 + 3x^2$
13	$w_6(x) = x^{10} + 4x^7 + 6x^4 + 2x$
24	$w_7(x) = x^{12} + 5x^9 + 10x^6 + 7x^3 + 1$

Remark 1.4 (cf. [18]). Let R be the ring of integers ($R = \mathbb{Z}$). Then we have

p	q	r	a	b	w_n is sequence of numbers
1	1	1	0	1	the Tribonacci number
1	1	1	1	1	the Tricobsthal number

Remark 1.5 (cf. [18]). Let R be the polynomial ring of integers ($R = \mathbb{Z}[x]$). Then we have

p	q	r	a	b	w_n is sequence of polynomials
x^2	x	1	0	1	the Tribonacci polynomial
1	x	x^2	1	1	the Tricobsthal polynomial

Let R be an integral domain. Let M be an $n \times n$ matrix with entries from R and let $\text{Adj}(M)$ be the adjugate of M . Then it is well-known that $M\text{Adj}(M) = \det(M)I$ and $\det(M)$ is invertible in R iff M is an invertible matrix (cf. [21]).

To benefit from the methods in linear algebra, we use the matrix $W(n)$ on sequences in [18] as follows:

$$W(n) = \begin{bmatrix} w_{n+1} & w_n & w_{n-1} \\ w_n & w_{n-1} & w_{n-2} \\ w_{n-1} & w_{n-2} & w_{n-3} \end{bmatrix}.$$

Lemma 1.6 (cf. [18]). For any integer n , we have the following relation

$$K.W(n - 1) = W(n),$$

where $K = \begin{bmatrix} p & q & r \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$.

In this article, some generalizations of the results in the literature are obtained by using the sequences over an integral domain with the help of matrix method. Then the generalizations for the d’Ocagne identity, the Honsberger’s formula, the Cassini’s identity and the Catalan’s identity are obtained with the help of them.

2. Tribonacci sequences by matrix method

Theorem 2.1. For integers n and m , the following hold:

- i) $W(n) = pW(n - 1) + qW(n - 2) + rW(n - 3)$.
- ii) $W(n + m) = K^n.W(m)$.

Proof. i) By Definition 1.1, it is clear.

ii) By Lemma 1.6, we have that $W(n) = K^n.W(0)$ for an integer n . Then it follows

$$W(n + m) = K^{n+m}.W(0) = K^n.K^m.W(0) = K^n.W(m)$$

□

When $w_0 = 0$ and $w_1 = 1$, we have a specific sequence (matrix) and so use the notation k_n ($K(n)$) instead of w_n ($W(n)$), respectively, i.e.

$$K(n) = \begin{bmatrix} k_{n+1} & k_n & k_{n-1} \\ k_n & k_{n-1} & k_{n-2} \\ k_{n-1} & k_{n-2} & k_{n-3} \end{bmatrix} \quad \text{and} \quad K(0) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & \frac{1}{r} \\ 0 & \frac{1}{r} & -\frac{q}{r^2} \end{bmatrix}.$$

Let us define the matrix $K_A(n)$ as $\begin{bmatrix} k_{n+1} & qk_n & 0 \\ k_n & qk_{n-1} & 0 \\ k_{n-1} & qk_{n-2} & 0 \end{bmatrix}$ and the matrix $K_B(n)$ as $\begin{bmatrix} 0 & rk_{n-1} & rk_n \\ 0 & rk_{n-2} & rk_{n-1} \\ 0 & rk_{n-3} & rk_{n-2} \end{bmatrix}$.

Theorem 2.2. For an integer n , we have the following:

- i) $K_A(n + 1) = K.(K_A(n))$.
- ii) $K_B(n + 1) = K.(K_B(n))$.
- iii) $K^n = K_A(n) + K_B(n)$.

Proof. i)

$$\begin{aligned} K.(K_A(n)) &= \begin{bmatrix} p & q & r \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} k_{n+1} & qk_n & 0 \\ k_n & qk_{n-1} & 0 \\ k_{n-1} & qk_{n-2} & 0 \end{bmatrix} \\ &= \begin{bmatrix} k_{n+2} & qk_{n+1} & 0 \\ k_{n+1} & qk_n & 0 \\ k_n & qk_{n-1} & 0 \end{bmatrix} \\ &= K_A(n + 1). \end{aligned}$$

ii)

$$\begin{aligned} K.(K_B(n)) &= \begin{bmatrix} p & q & r \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 0 & rk_{n-1} & rk_n \\ 0 & rk_{n-2} & rk_{n-1} \\ 0 & rk_{n-3} & rk_{n-2} \end{bmatrix} \\ &= \begin{bmatrix} 0 & rk_n & rk_{n+1} \\ 0 & rk_{n-1} & rk_n \\ 0 & rk_{n-2} & rk_{n-1} \end{bmatrix} \\ &= K_B(n+1). \end{aligned}$$

iii) By the induction method, we have completed the proof. □

Theorem 2.3. For integers n and m , we have the following for the sequence $\{w_n\}$.

1) Honsberger's formula:

$$w_{m+n+1} = k_{n+1}w_{m+1} + w_m(qk_n + rk_{n-1}) + rk_nw_{m-1}.$$

2) Catalan's identity: $\det(W(n)) = r^m \det(W(n-m))$.

3) Cassini's or Simson's identity:

$$2w_nw_{n-1}w_{n-2} + w_{n+1}w_{n-1}w_{n-3} - w_{n-1}^3 - w_{n+1}w_{n-2}^2 - w_{n-3}w_n^2 = r^n (-w_{-3}w_0^2 - w_1w_{-2}^2).$$

4)

$$2k_nk_{n-1}k_{n-2} + k_{n+1}k_{n-1}k_{n-3} - k_{n-1}^3 - k_{n+1}k_{n-2}^2 - k_{n-3}k_n^2 = -r^{n-2}.$$

5)

$$w_{n+1} = k_{n+1}b + a(qk_n + rk_{n-1}).$$

Proof. By Theorem 2.1, we prove the following.

1) We have

$$W(n+m) = K^n.W(m).$$

By the matrix equality, we get that

$$w_{m+n+1} = k_{n+1}w_{m+1} + w_m(qk_n + rk_{n-1}) + rk_nw_{m-1}.$$

2) We have $W(n) = K^m.W(n-m)$, for any integers m and n . Then, by the determinant, it follows that

$$\det(W(n)) = r^m \det(W(n-m)).$$

3) We have $W(n) = K^n.W(0)$. Then, by the determinant, it follows that

$$\begin{aligned} &2w_nw_{n-1}w_{n-2} + w_{n+1}w_{n-1}w_{n-3} - w_{n-1}^3 - w_{n+1}w_{n-2}^2 - w_{n-3}w_n^2 \\ &= r^n (-w_{-3}w_0^2 + 2w_0w_{-1}w_{-2} - w_{-1}^3 + w_1w_{-3}w_{-1} - w_1w_{-2}^2). \end{aligned}$$

4) We have

$$\begin{aligned} 2k_nk_{n-1}k_{n-2} + k_{n+1}k_{n-1}k_{n-3} - k_{n-1}^3 - k_{n+1}k_{n-2}^2 - k_{n-3}k_n^2 &= r^n (-k_{-2}^2) \\ &= -r^{n-2}. \end{aligned}$$

5) It is clear from (i). □

Corollary 2.4. For any integers n and m , we have the following equalities for the n -th Tribonacci numbers T_n and the n -th Tricobsthal numbers J_n .

1)

$$\begin{aligned} T_{m+n} &= T_{n+1}T_m + T_{m-1}(T_n + T_{n-1}) + T_nT_{m-2}. \\ J_{m+n} &= T_{n+1}J_m + J_{m-1}(T_n + T_{n-1}) + T_nJ_{m-2}. \end{aligned}$$

2)

$$\begin{aligned} -1 &= 2J_nJ_{n-1}J_{n-2} + J_{n+1}J_{n-1}J_{n-3} - J_{n-1}^3 - J_{n+1}J_{n-2}^2 - J_{n-3}J_n^2 \\ &= 2T_nT_{n-1}T_{n-2} + T_{n+1}T_{n-1}T_{n-3} - T_{n-1}^3 - T_{n+1}T_{n-2}^2 - T_{n-3}T_n^2 \end{aligned}$$

3)

$$J_{n+1} = T_{n+1} + T_n + T_{n-1}.$$

The sequence in Definition 1.1 equals the sequence defined in [1] when $r = 0$ and so the following equalities are obtained from [1].

Corollary 2.5 (cf. [1]). For any integers n and m , we have the following equalities.

1) Catalan's identity:

$$(-q)^m w_{n-m}^2 - w_n^2 = (-q)^m w_{n-m-1}w_{n-m+1} - w_{n-1}w_{n+1}.$$

2) Cassini's or Simson's identity:

$$w_{n-1}w_{n+1} - w_n^2 = (-q)^n (w_1^2 - qw_0^2 - pw_0w_1).$$

3) $k_{n-1}k_{n+1} - k_n^2 = (-q)^{n-1}$.

4) Honsberger's formula: $w_{m+n} = k_n w_{m+1} + qk_{n-1}w_m$.

5) d'Ocagne's identity: $(-q)^n w_m = k_{n+1}w_{m+n} - k_n w_{m+n+1}$.

Theorem 2.6. For any integer n , we have

$$w_{-n} = \frac{1}{r^{n-1}} \left(\begin{vmatrix} k_{n+1} & k_n \\ k_{n-1} & k_{n-2} \end{vmatrix} a + (-q)^{n-2} b \right).$$

Proof. To obtain w_{-n} , we compute the $(2, 1)$ th entry in the matrix $W(-n)$ by Theorem 2.1 as follows:

$$\begin{aligned} w_{-n} &= \frac{1}{r^n} \left[- \begin{vmatrix} k_n & rk_{n-1} \\ k_{n-1} & rk_{n-2} \end{vmatrix} \begin{vmatrix} k_{n+1} & rk_n \\ k_{n-1} & rk_{n-2} \end{vmatrix} - \begin{vmatrix} k_{n+1} & rk_n \\ k_n & rk_{n-1} \end{vmatrix} \right] \begin{bmatrix} w_1 \\ w_0 \\ w_{-1} \end{bmatrix} \\ &= \frac{1}{r^{n-1}} \left(- \begin{vmatrix} k_n & k_{n-1} \\ k_{n-1} & k_{n-2} \end{vmatrix} b + \begin{vmatrix} k_{n+1} & k_n \\ k_{n-1} & k_{n-2} \end{vmatrix} a \right) \end{aligned}$$

Then we have $w_{-n} = \frac{1}{r^{n-1}} \left(\begin{vmatrix} k_{n+1} & k_n \\ k_{n-1} & k_{n-2} \end{vmatrix} a - \begin{vmatrix} k_n & k_{n-1} \\ k_{n-1} & k_{n-2} \end{vmatrix} b \right)$.

From (3) in Corollary 2.5, we have that $k_{n-1}k_{n+1} - k_n^2 = (-q)^{n-1}$. Thus

$$w_{-n} = \frac{1}{r^{n-1}} \left(\begin{vmatrix} k_{n+1} & k_n \\ k_{n-1} & k_{n-2} \end{vmatrix} a + (-q)^{n-2} b \right).$$

□

Corollary 2.7. Let n and m be integers. Then we have

$$w_{m-n} = \frac{1}{r^{n-1}} \left(\begin{vmatrix} k_{n+1} & k_n \\ k_{n-1} & k_{n-2} \end{vmatrix} w_m - (-q)^{n-2} (w_{m+1} + qw_{m-1}) \right).$$

Let us compute w_{-2} by using Theorem 2.6.

Example 2.8. Since $k_0 = 0, k_1 = 1, k_2 = p, k_3 = p^2 + q$, w_{-2} can be computed as follows:

$$\begin{aligned} w_{-2} &= \frac{1}{r} \left(\begin{vmatrix} k_3 & k_2 \\ k_1 & k_0 \end{vmatrix} a + (-q)^0 b \right) \\ &= \frac{b - pa}{r}. \end{aligned}$$

Theorem 2.9. Let $t^3 + pt^2 + qt + r = (t - \alpha)(t - \beta)(t - \delta)$ for some $\alpha, \beta, \gamma \in R$, which are different from each others. Then for any integer n , we have

$$\begin{aligned} k_n &= \frac{\alpha^{n+1}}{(\alpha - \gamma)(\alpha - \beta)} - \frac{\beta^{n+1}}{(\beta - \gamma)(\alpha - \beta)} + \frac{\gamma^{n+1}}{(\beta - \gamma)(\alpha - \gamma)} \\ w_n &= w_1 k_n - w_0 \left(\frac{\alpha^{n+1}(\beta + \gamma)}{(\alpha - \gamma)(\alpha - \beta)} - \frac{\beta^{n+1}(\alpha + \gamma)}{(\beta - \gamma)(\alpha - \beta)} + \frac{\gamma^{n+1}(\alpha + \beta)}{(\beta - \gamma)(\alpha - \gamma)} \right). \end{aligned}$$

Proof. Let α, β, γ be elements of R , which are different from each others and $t^3 + pt^2 + qt + r = (t - \alpha)(t - \beta)(t - \delta)$. Let P and D be the matrices as follows:

$$P = \begin{bmatrix} \alpha^2 & \beta^2 & \gamma^2 \\ \alpha & \beta & \gamma \\ 1 & 1 & 1 \end{bmatrix} \quad \text{and} \quad D = \begin{bmatrix} \alpha & 0 & 0 \\ 0 & \beta & 0 \\ 0 & 0 & \gamma \end{bmatrix}.$$

Then we get that $\det(P) = (\beta - \gamma)(\alpha - \gamma)(\alpha - \beta)$.

Moreover, it is clear that $\det(P)K = (PD)(\text{Adj}(P))$ and thus we have $\det(P)K^n = (PD^n)(\text{Adj}(P))$. It follows that

$$\det(P)W(n) = \det(P)K^n W(0) = (PD^n)(\text{Adj}(P))W(0).$$

By the matrix equality, we get that

$$\begin{aligned} \det(P)w_{n+1} &= w_1 (\alpha^2 \alpha^n (\beta - \gamma) - \beta^2 \beta^n (\alpha - \gamma) + \gamma^2 \gamma^n (\alpha - \beta)) \\ &\quad - w_0 (\alpha^2 \alpha^n (\beta^2 - \gamma^2) - \beta^2 \beta^n (\alpha^2 - \gamma^2) + \gamma^2 \gamma^n (\alpha^2 - \beta^2)) \\ &\quad - w_{-1} (\alpha^2 \alpha^n (\beta \gamma^2 - \beta^2 \gamma) - \beta^2 \beta^n (\alpha \gamma^2 - \alpha^2 \gamma) + \gamma^2 \gamma^n (\alpha \beta^2 - \alpha^2 \beta)). \end{aligned}$$

By the definition of the sequence k_n and since $k_0 = 0, k_1 = 1, k_{-1} = w_{-1} = 0$, we have that

$$\det(P)k_{n+1} = (\alpha^2 \alpha^n (\beta - \gamma) - \beta^2 \beta^n (\alpha - \gamma) + \gamma^2 \gamma^n (\alpha - \beta))$$

and so

$$\begin{aligned} k_{n+1} &= \frac{(\alpha^2 \alpha^n (\beta - \gamma) - \beta^2 \beta^n (\alpha - \gamma) + \gamma^2 \gamma^n (\alpha - \beta))}{(\beta - \gamma)(\alpha - \gamma)(\alpha - \beta)} \\ &= \frac{\alpha^{n+2}}{(\alpha - \gamma)(\alpha - \beta)} - \frac{\beta^{n+2}}{(\beta - \gamma)(\alpha - \beta)} + \frac{\gamma^{n+2}}{(\beta - \gamma)(\alpha - \gamma)}. \end{aligned}$$

Again we focus on the above equation

$$\begin{aligned} w_{n+1} &= w_1 \frac{(\alpha^2 \alpha^n (\beta - \gamma) - \beta^2 \beta^n (\alpha - \gamma) + \gamma^2 \gamma^n (\alpha - \beta))}{(\beta - \gamma)(\alpha - \gamma)(\alpha - \beta)} \\ &\quad - w_0 \frac{(\alpha^2 \alpha^n (\beta^2 - \gamma^2) - \beta^2 \beta^n (\alpha^2 - \gamma^2) + \gamma^2 \gamma^n (\alpha^2 - \beta^2))}{(\beta - \gamma)(\alpha - \gamma)(\alpha - \beta)} \\ &\quad - w_{-1} \frac{(\alpha^2 \alpha^n (\beta \gamma^2 - \beta^2 \gamma) - \beta^2 \beta^n (\alpha \gamma^2 - \alpha^2 \gamma) + \gamma^2 \gamma^n (\alpha \beta^2 - \alpha^2 \beta))}{(\beta - \gamma)(\alpha - \gamma)(\alpha - \beta)} \\ &= w_1 k_{n+1} - w_0 \left(\frac{\alpha^{n+2}(\beta + \gamma)}{(\alpha - \gamma)(\alpha - \beta)} - \frac{\beta^{n+2}(\alpha + \gamma)}{(\beta - \gamma)(\alpha - \beta)} + \frac{\gamma^{n+2}(\alpha + \beta)}{(\beta - \gamma)(\alpha - \gamma)} \right) - \alpha \beta \gamma w_{-1} k_n \\ &= a k_{n+1} - b \left(\frac{\alpha^{n+2}(\beta + \gamma)}{(\alpha - \gamma)(\alpha - \beta)} - \frac{\beta^{n+2}(\alpha + \gamma)}{(\beta - \gamma)(\alpha - \beta)} + \frac{\gamma^{n+2}(\alpha + \beta)}{(\beta - \gamma)(\alpha - \gamma)} \right). \end{aligned}$$

since $w_{-1} = 0$. Then the proof is completed. □

Acknowledgement

The authors thank the referees for valuable suggestions and comments.

References

- [1] M. Alkan, *The generalized Fibonacci sequences on an integral domain*, Montes Taurus J. Pure Appl. Math. **3** (2), 60–69, 2021.
- [2] G. E. Bergum and V. E. Hoggatt, *Sums and products for recurring sequences*, Fibonacci Quart. **13** (2), 115–120, 1975.
- [3] G. S. Cheon, H. Kim and L. W. Shapiro, *A generalization of Lucas polynomial sequence*, Discrete Appl. Math. **157** (5), 920–927, 2009.
- [4] E. Choi, *Modular Trinonacci numbers by matrix method*, J. Korean Soc. Math. Educ. Ser. B: Pure Appl. Math. **20** (3), 207–221, 2013.
- [5] S. Falcón, *On the k -Lucas numbers*, Int. J. Contemp Math. Sci. **6** (21-24), 1039–1050, 2011.
- [6] S. Falcón and Á. Plaza, *On k -Fibonacci sequences and polynomials and their derivatives*, Chaos, Solitons & Fractals **39**, 1005–1019, 2009.
- [7] A. F. Horadam, *Basic properties of a certain generalized sequence of numbers*, Fibonacci Quart. **3** (2), 161–176, 1965.
- [8] A. F. Horadam, *Tschebyscheff and other functions associated with the sequence $W_n(a, b; p, q)$* , Fibonacci Quart. **7** (1), 14–22, 1969.
- [9] A. F. Horadam, *A synthesis of certain polynomial sequences*, Applications of Fibonacci numbers, Kluwer Academic Publishers, 215–229, Dordrecht, 1996.
- [10] L. C. Hsu and P. J. S. Shiue, *Cycle indicators and special functions*, Ann. Comb. **5** (2), 179–196, 2001.
- [11] M. Janjic, *Hessenberg matrices and integer sequences*, J. Integer Seq. **13** (7), Article ID: 10.7.8, 2010.
- [12] T. Koshy, *Fibonacci and Lucas numbers with applications*, Wiley-Interscience Publication, New York, 2001.
- [13] G. Lee and M. Asci, *Some properties of the (p, q) -Fibonacci and (p, q) -Lucas polynomials*, J. Appl. Math., Article ID: 264842, 2012.
- [14] H. Merzouk, A. Boussayoud and M. Chelgham, *Generating functions of generalized Tribonacci and Tricobsthal polynomials*, Montes Taurus J. Pure Appl. Math. **2** (2), 7–37, 2020.
- [15] A. Nalli and P. Haukkanen, *On generalized Fibonacci and Lucas polynomials*, Chaos, Solitons & Fractals **42** (5), 3179–3186, 2009.
- [16] G. Ozdemir and Y. Simsek, *Generating functions for two-variable polynomials related to a family of Fibonacci type polynomials and numbers*, Filomat **30** (4), 969–975, 2016.
- [17] G. Ozdemir, Y. Simsek and G. V. Milovanović, *Generating functions for special polynomials and numbers including Apostol-type and Humbert-type polynomials*, Mediterr. J. Math. **14** (3), 1–17, 2017.
- [18] Ö. Öneş and M. Alkan, *On the generalized Tribonacci sequences*, Proceedings Book of Micopam 2020-2021, ISBN: 978-625-00-0397-8, 39-42, 2021.
- [19] J. L. Ramírez, *Some properties of convolved k -Fibonacci numbers*, ISRN Combin. Article ID: 759641, 2013.
- [20] J. L. Ramírez, *On convolved generalized Fibonacci and Lucas polynomials*, Appl. Math. Comput. **229**, 208–213, 2014.
- [21] S. Roman, *Advanced Linear Algebra*, New York, Springer, 2008.
- [22] B. Rybolowicz and A. Tereszkiwicz, *Generalized Tricobsthal and Tribonacci Polynomials*, Appl. Math. Comput. **325**, 297–308, 2018.
- [23] H. Tian-Xiao and J. S. Peter, *On sequences of numbers and polynomials defined by linear recurrence relations of order 2*, Int. J. Math. Sci. **21**, Article ID: 709386.356, 2009.
- [24] M. E. Waddill and L. Sacks, *Another generalized Fibonacci sequence*, Fibonacci Quart. **5** (3), 209–222, 1967.
- [25] W. Weiping and W. Hui, *Generalized Humbert polynomials via generalized Fibonacci polynomials*, Appl. Math. Comput. **307**, 204–216, 2017.
- [26] J. Wang, *Some new results for the (p, q) -Fibonacci and Lucas polynomials*, Adv. Differ. Equ. **2014**, Article ID: 64, 2014.
- [27] F. Z. Zhao and T. Wang, *Some identities involving the powers of the generalized Fibonacci numbers*, Fibonacci Quart. **41** (1), 7–12, 2003.